

**ARIZONA GAME AND FISH DEPARTMENT  
HERITAGE DATA MANAGEMENT SYSTEM**

**Animal Abstract**

**Element Code:** ABNSB08041

**Data Sensitivity:** Yes

**CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE**

**NAME:** *Glaucidium brasilianum cactorum* van Rossem

**COMMON NAME:** Cactus Ferruginous Pygmy-Owl

**SYNONYMS:** *Strix brasiliana* Gmelin

**FAMILY:** Strigidae

**AUTHOR, PLACE OF PUBLICATION:** Van Rossem. 1937. Proc. Biol. Soc. Washington, 50, Feb. 23, 1937. p.27.

**TYPE LOCALITY:** Between Guaymas and Empalme, Sonora.

**TYPE SPECIMEN:** 30225 California Institute of Technology (original number 12940).

**TAXONOMIC UNIQUENESS:** The subspecies *cactorum* is 1 of 3 subspecies of *Glaucidium brasilianum*, and is the only subspecies that occurs in Arizona. *G. b. ridgwayi* occurs throughout southern Mexico and Central America, and *G. b. brasilianum* is found throughout South America.

**DESCRIPTION:** A small size owl weighing 2.3-3.1 ounces and measuring 5.8-7.2 in (14.73-18.3 cm) in length, with a 14.3-16.0 in (36.32-40.64 cm) wingspan (Monson *in* Glinski 1998). Females average slightly larger than males. The back and upper coverts are plain light brown, with occasional conspicuous white spots on the upper wing coverts and small, regularly spaced whitish spots on the outer primaries. Both the tail and the flight feathers are dusky barred, the dusky and light bars of the tail evenly spaced. The head and nape are a somewhat paler brown than the back, and have linear whitish streaks. Each side of the nape is decorated with a black blot bordered in white that resembles an eye. The face has a white V and disk. The breast and flanks are heavily streaked with dark reddish brown to blackish marks; sometimes the streaks merge and become almost solid reddish brown. The tarsi are feathered. The eyes are yellow, and the bill is greenish-yellow. (Monson *in* Glinski 1998). Both sexes and immature birds have similar plumage.

**AIDS TO IDENTIFICATION:** Similar to Northern Pygmy-Owl (*Glaucidium gnoma*). Rufous crown streaked with white. Tail has 7-8 light brown (not white) bars. Their call consists of a “put-put-put” note with the “sharp penetrating quality of the whistling note of the cardinal; the rate, however, being about 150 ‘puts’ per minute.” Early in season, the male may call “nearly all night and much of the day.” (Brandt 1951).

**ILLUSTRATIONS:**

Color drawing of species (Monson *in* Glinski 1998: plate 36)

**TOTAL RANGE:** Southern Arizona and southern Texas, south to Guerrero, Nuevo Leon, and Tamaulipas, Mexico.

**RANGE WITHIN ARIZONA:** The only recent records are from Organ Pipe Cactus National Monument, near Ajo, and suburban Tucson. Historically, fairly common in mesquite bosques throughout central southern Arizona. They have been found south of Tortolita Mts., west of Tortilla Mts., Rincon Mts., Pajarito Mts., Puerto Blanco Mts., Ajo Mts., Santa Catalina Mts., Santa Rita Mts., Tucson area, Gila River near Bonita Creek and San Francisco River, San Pedro River near Dudleyville, and Sonoyta Creek.

### **SPECIES BIOLOGY AND POPULATION TRENDS**

**BIOLOGY:** A nonsocial owl that is active mostly at dawn and dusk. This diurnal habit helps separate it from other small owls that dislike light. Vocalizations indicate breeding activity, usually September to April (Monson *in* Glinski 1998; September to March in Millsap and Johnson *in press*). It is normally silent in the summer. The principal vocalization is a rapid, monotonous repeated and shrill hooting (Monson *in* Glinski 1998). The hoots are uttered at the rate of about 2 per second, or 120-150 per minute, and may be repeated as many as 100 times without pause (Monson *in* Glinski 1998). The flight is quick and direct, generally flying short distances from one tree or bush to another. When perching, it usually sits in a leafy paloverde mesquite, or perhaps a cottonwood or willow. The best field identification is its small size and long reddish-barred tail, which is often nervously wagged or twitched. It is unknown whether they remain on their home territory throughout the year.

**REPRODUCTION:** Nothing is known about the courtship behavior (Monson *in* Glinski 1998). They nest in woodpecker or natural cavities in broadleaf riparian trees or saguaro cacti. Nests usually 10-20 feet up but may be as high as 40 feet and close to the river. No lining material used. Laying commences by 20 April. Three to 4 white, unmarked eggs (1.1 inches long) are laid in the bottom of the cavity. Possibly monogamous mating. Twenty-eight day incubation period, mostly by the female. Young are immobile, downy, and have eyes closed after hatching. Both parents feed the young; the male will also feed the female. Hatching is synchronous but siblings compete for food resulting in size differences. Young fledge in 27-30 days.

**FOOD HABITS:** A daytime predator that's diet is not well known. It forages in microphyllous woodland, especially well developed mesquite bosque, but also desert washes with mature blue paloverde, mesquite, and ironwood. Small birds, insects (e.g. crickets, scorpions, and caterpillars), lizards, and mammals thought to be their main diet. Prey is snatched from the ground in talons, after a gliding descent from a perch.

**HABITAT:** In Arizona according to Monson (*in* Glinski 1998), "it has occurred in streamside cottonwoods and willows and adjacent mesquite bosques, usually with saguaros on nearby

slopes. Less often it has been found along dry washes where large mesquite, paloverde, ironwood, and saguaro thrive.”

**ELEVATION:** 1,300 - 4,000 ft (397 - 1,220 m).

**PLANT COMMUNITY:** Sonoran riparian deciduous woodland, within Arizona upland subdivision and Sonoran desertscrub.

**POPULATION TRENDS:** Drastic declines in both range and abundance. Sporadic breeding has been documented in recent years in suburban Tucson. Before 1950, they were a fairly common to common resident in low elevation riparian mesquite woodlands ranging north to Phoenix, northwest to Salt-Gila River confluence, west to Cabeza Prieta Tanks, and east to at least Superior (probably to upper Gila River near Safford). They are declining in northern Sonora where they now are absent from many locations described by van Rossem.

## **SPECIES PROTECTION AND CONSERVATION**

<b>ENDANGERED SPECIES ACT STATUS:</b>	SC (USDI, FWS 2006) [LE (USDI, FWS 1997); Critical Habitat Proposed 11/27/2002, AZ Pop.] [PE USDI, FWS 1996] [C1 USDI, FWS 1991]
<b>STATE STATUS:</b>	1B (AGFD SWAP 2012) [WSC, AGFD, WSCA in prep] [State Endangered (species level) AGFD, TNW 1988]
<b>OTHER STATUS:</b>	Bureau of Land Management Sensitive (USDI, BLM AZ 2008, 2010) Forest Service Sensitive (USDA, FS Region 3 1999, 2007, 2013) Texas Parks and Wildlife Department – Threatened. None. (NORMA Oficial Mexicana NOM- 059-SEMARNAT-2010). [A (Diario Oficial de la Federacion 1994), listed as full species.]

**MANAGEMENT FACTORS:** **Likely threats:** degradation and loss of habitat and urban development in saguaro-ironwood forests (near Tucson). **Management needs:** refine breeding survey protocols and conduct surveys; determine habitat needs; manage grazing and other land uses to maintain and enhance important occupied or potentially occupied habitats. (AGFD in prep).

**PROTECTIVE MEASURES TAKEN:**

**SUGGESTED PROJECTS:** Continued statewide status surveys. Refinement of detection protocol, determination of habitat requirements needed.

**LAND MANAGEMENT/OWNERSHIP:** BIA - Tohono O'odham Nation; BLM - Tucson Field Office; FWS - Buenos Aires and Cabeza Prieta National Wildlife Refuges; NPS - Organ Pipe Cactus National Monument and Saguaro National Park; USFS - Coronado and Tonto National Forests; State Land Department; Private.

## **SOURCES OF FURTHER INFORMATION**

### **REFERENCES:**

- Abbate, D., Ditty, A., Richardson, S. and Olding, R. 1996. Cactus ferruginous pygmy-owl surveys and nest monitoring in the Tucson Basin area, Arizona - 1996. Arizona Game and Fish Department. Heritage Grant Report #U95503. 35 pages.
- Arizona Game and Fish Department. 1988. Threatened Native Wildlife in Arizona. p. 15.
- Arizona Game and Fish Department (AGFD). In prep. Wildlife of special concern in Arizona. Arizona Game and Fish Department Publication. Phoenix, Arizona. 32 pp.
- Arizona Game and Fish Department. 2012. Arizona's State Wildlife Action Plan 2012-2022. Phoenix, AZ.
- Brandt, H. 1951. Arizona and its bird life. The Bird Research Foundation, Cleveland, Ohio. p.653.
- Collins, M.D. and Corman, T.E. 1995. Cactus ferruginous pygmy-owl surveys in Arizona: 1993-1994 season. Arizona Game and Fish Department. Technical Report 37. 17 pages.
- Felley, D.L. and Corman, T.E. 1993. Spring 1993 cactus ferruginous pygmy-owl surveys in Arizona. Arizona Game and Fish Department. Technical Report 36. 16 pages.
- Hunter, W.C. Status of the cactus ferruginous pygmy-owl (*Glaucidium brasilianum cactorum*) in the United States and northern Mexico. USDI Fish and Wildlife Service, Phoenix, Arizona. 13 pp.
- Lesh, T.D. and Corman, T.E. 1995. Cactus ferruginous pygmy-owl surveys in Arizona: 1993-1995. Arizona Game and Fish Department. Technical Report 76. 23 pages.
- Millsap, B.A. and R.R. Johnson. 1988. In R.L. Glinski et al. (eds.). Proceedings of the Southwest Raptor Management Symposium and Workshop. National Wildlife Federation, Washington, D.C. 395 pp.
- Monson, G. Ferruginous Pygmy - Owl: *Glaucidium brasilianum*. pp. 159-161. In Glinski, R.L. (Ed.). 1998. The Raptors of Arizona. The University of Arizona Press and the Arizona Game and Fish Department.
- Proyecto de Norma Oficial Mexicana. 2000. PROY-NOM-059-ECOL-2000.
- Secretaría de Medio Ambiente y Recursos Naturales. 2010. NORMA Oficial Mexicana NOM-059-SEMARNAT-2010, Protección ambiental-Especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en riesgo.
- Terres, J.K. 1980. The Audubon Society encyclopedia of North American birds. Alfred A. Knopf, New York. p.665.

- USDA, Forest Service Region 3. 1999(2000). Regional Forester's Sensitive Species List.
- USDA, Forest Service Region 3. 2007. Regional Forester's List of Sensitive Animals.
- USDA, Forest Service Region 3. 2013. Regional Forester's List of Sensitive Animals.
- USDI, Bureau of Land Management. 2008. Arizona BLM Sensitive Species List.
- USDI, Bureau of Land Management. 2010. Arizona BLM Sensitive Species List.
- USDI, Fish and Wildlife Service. 1991. Endangered and Threatened Wildlife and Plants; Animal Candidate Review for Listing as Endangered or Threatened Species, Proposed Rule. Federal Register 56(225):58811.
- USDI, Fish and Wildlife Service. 1994. Endangered and Threatened Wildlife and Plants; Animal Candidate Review for Listing as Endangered or Threatened Species; Proposed Rule. Federal Register 59(219):58991.
- USDI, Fish and Wildlife Service. 1994. Proposed Rule to list the cactus ferruginous pygmy-owl as endangered with critical habitat in Arizona and threatened in Texas. Federal Register 59:237 63975-63986.
- USDI, Fish and Wildlife Service. 1996. Endangered and Threatened Wildlife and Plants: Review of Plant and Animal Taxa that are Candidates for Listing as Endangered or Threatened Species. Federal Register 61(40):7596-7613.
- USDI, Fish and Wildlife Service. 1997. Endangered and Threatened Wildlife and Plants; Determination of Endangered Status for the Cactus Ferruginous Pygmy-Owl in Arizona; Final Rule Federal Register 62(46):10730-10747.
- USDI, Fish and Wildlife Service. 1999. Federal Register 64(132):37419-37440.
- USDI, Fish and Wildlife Service. 2002. Endangered and Threatened Wildlife and Plants; Designation of Critical Habitat for the Arizona Distinct Population Segment of the Cactus Ferruginous Pygmy-owl (*Glaucidium brasilianum cactorum*); Proposed rule; notice of availability. Federal Register 67(229):71032-71064.
- USDI, Fish and Wildlife Service. 2006. Endangered and Threatened Wildlife and Plants; Final Rule to Remove the Arizona Distinct Population Segment of the Cactus Ferruginous Pygmy-owl (*Glaucidium brasilianum cactorum*) From the Federal List of Endangered and Threatened Wildlife; Withdrawal of the Proposed Rule to Designate Critical Habitat; Removal of Federally Designated Critical Habitat; Final Rule. Federal Register 71(72): 19452-19458.

**MAJOR KNOWLEDGEABLE INDIVIDUALS:**

C. Hunter - US Fish and Wildlife Service  
Roy Johnson - National Park Service, University of Arizona, Tucson.  
Brian Millsap - Florida Game and Fresh Water Fish Commission, Lakeland, Florida.  
S. Russell - University of Arizona, Tucson.  
Troy Corman - Arizona Game and Fish Department, Phoenix.  
Scott Richardson – USFWS  
Marty Teagle - USFWS.

**ADDITIONAL INFORMATION:**

“The name *Glaucidium* is derived from the Greek *glaux*, which implies that this bird is an owl because it has glaring eyes (Choate 1985); *brasilianum* refers to Brazil, where this small diurnal owl was discovered. The subspecific epithet for the race occurring in Arizona,

*cactorum*, alludes to cactus, a vegetation that it frequently associates with in the northern portion of its range. The term *ferruginous* refers to the rusty red tail and upper side of the outer vanes of the flight feathers.” (Monson *in* Glinski 1998).

**Revised:** 1992-01-17 (JGH)  
1995-03-21 (LZW)  
1997-03-05 (SMS)  
1998-03-02 (SSS)  
2001-06-08 (SMS)

To the user of this abstract: you may use the entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.